

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Vascular - angina (chest pain) (Q1)							
-	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	230	165	56	45	48	544
	%	73.24%	85.49%	84.84%	84.90%	40.67%	73.11%
- Recent/active	N	10	3	--	3	3	19
	%	3.18%	1.55%	--	5.66%	2.54%	2.55%
- Remote/inactive	N	14	17	7	4	4	46
	%	4.45%	8.80%	10.60%	7.54%	3.38%	6.18%
- Not determined	N	2	1	1	1	--	5
	%	0.63%	0.51%	1.51%	1.88%	--	0.67%
Vascular - heart attack / cardiac arrest (Q2)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	132	182	59	49	53	475
	%	42.03%	94.30%	89.39%	92.45%	44.91%	63.84%
- Recent/active	N	3	--	--	1	1	5
	%	0.95%	--	--	1.88%	0.84%	0.67%
- Remote/inactive	N	18	4	5	3	1	31
	%	5.73%	2.07%	7.57%	5.66%	0.84%	4.16%
Vascular - hypertension (high blood pressure) (Q3)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	63	113	26	22	22	246
	%	20.06%	58.54%	39.39%	41.50%	18.64%	33.06%
- Recent/active	N	78	64	33	26	28	229
	%	24.84%	33.16%	50.00%	49.05%	23.72%	30.77%
- Remote/inactive	N	12	8	5	2	4	31
	%	3.82%	4.14%	7.57%	3.77%	3.38%	4.16%
- Not determined	N	--	1	--	3	1	5
	%	--	0.51%	--	5.66%	0.84%	0.67%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
		314	193	66	53	118	744	
Vascular - hyperlipidemia (high cholesterol, LDL, or triglycerides (Q4))								
	N	161	7	2	--	63	233	
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%	
- Absent		N	58	79	24	17	16	194
		%	18.47%	40.93%	36.36%	32.07%	13.55%	26.07%
- Recent/active		N	86	96	34	34	32	282
		%	27.38%	49.74%	51.51%	64.15%	27.11%	37.90%
- Remote/inactive		N	9	9	6	2	6	32
		%	2.86%	4.66%	9.09%	3.77%	5.08%	4.30%
- Not determined		N	--	2	--	--	1	3
		%	--	1.03%	--	--	0.84%	0.40%
Vascular - stroke (Q5)								
	N	160	7	2	--	63	232	
	%	50.95%	3.62%	3.03%	--	53.38%	31.18%	
- Absent		N	140	178	58	52	47	475
		%	44.58%	92.22%	87.87%	98.11%	39.83%	63.84%
- Recent/active		N	2	3	--	--	2	7
		%	0.63%	1.55%	--	--	1.69%	0.94%
- Remote/inactive		N	10	5	6	1	6	28
		%	3.18%	2.59%	9.09%	1.88%	5.08%	3.76%
- Not determined		N	2	--	--	--	--	2
		%	0.63%	--	--	--	--	0.26%
Vascular - TIA (stroke like event < 24 hours) (Q6)								
	N	161	7	2	--	63	233	
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%	
- Absent		N	138	173	59	44	44	458
		%	43.94%	89.63%	89.39%	83.01%	37.28%	61.55%
- Recent/active		N	3	1	1	3	2	10
		%	0.95%	0.51%	1.51%	5.66%	1.69%	1.34%
- Remote/inactive		N	12	9	3	4	5	33
		%	3.82%	4.66%	4.54%	7.54%	4.23%	4.43%

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
Vascular - TIA (stroke like event < 24 hours) (Q6)								
- Not determined		N	--	3	1	2	4	10
		%	--	1.55%	1.51%	3.77%	3.38%	1.34%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
		314	193	66	53	118	744	
Vascular - atrial fibrillation (certain heart rhythm problem) (Q7)								
	N	161	7	2	--	63	233	
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%	
- Absent		N	126	173	55	43	52	449
		%	40.12%	89.63%	83.33%	81.13%	44.06%	60.34%
- Recent/active		N	21	7	6	9	2	45
		%	6.68%	3.62%	9.09%	16.98%	1.69%	6.04%
- Remote/inactive		N	6	6	3	1	--	16
		%	1.91%	3.10%	4.54%	1.88%	--	2.15%
- Not determined		N	--	--	--	--	1	1
		%	--	--	--	--	0.84%	0.13%
Vascular - pacemaker (Q8)								
	N	161	7	2	--	63	233	
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%	
- Absent		N	139	180	61	51	55	486
		%	44.26%	93.26%	92.42%	96.22%	46.61%	65.32%
- Recent/active		N	14	6	2	2	--	24
		%	4.45%	3.10%	3.03%	3.77%	--	3.22%
- Remote/inactive		N	--	--	1	--	--	1
		%	--	--	1.51%	--	--	0.13%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Neurological - idiopathic Parkinson's disease (Q9)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	151	185	62	53	52	503
	%	48.08%	95.85%	93.93%	100.00%	44.06%	67.60%
- Recent/active	N	2	1	1	--	3	7
	%	0.63%	0.51%	1.51%	--	2.54%	0.94%
- Not determined	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%
Neurological - head Trauma (Q10)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	121	127	47	32	34	361
	%	38.53%	65.80%	71.21%	60.37%	28.81%	48.52%
- Recent/active	N	9	11	4	4	11	39
	%	2.86%	5.69%	6.06%	7.54%	9.32%	5.24%
- Remote/inactive	N	21	46	13	17	7	104
	%	6.68%	23.83%	19.69%	32.07%	5.93%	13.97%
- Not determined	N	2	2	--	--	3	7
	%	0.63%	1.03%	--	--	2.54%	0.94%
Neurological - seizure disorder (Q11)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	150	184	61	51	55	501
	%	47.77%	95.33%	92.42%	96.22%	46.61%	67.33%
- Recent/active	N	--	--	1	1	--	2
	%	--	--	1.51%	1.88%	--	0.26%
- Remote/inactive	N	2	2	1	1	--	6
	%	0.63%	1.03%	1.51%	1.88%	--	0.80%

(Continued)

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Neurological - seizure disorder (Q11)							
- Not determined	N	1	--	1	--	--	2
	%	0.31%	--	1.51%	--	--	0.26%
Neurological - incontinence (Q12)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	91	131	38	28	23	311
	%	28.98%	67.87%	57.57%	52.83%	19.49%	41.80%
- Recent/active	N	51	40	21	21	29	162
	%	16.24%	20.72%	31.81%	39.62%	24.57%	21.77%
- Remote/inactive	N	10	15	4	4	3	36
	%	3.18%	7.77%	6.06%	7.54%	2.54%	4.83%
- Not determined	N	1	--	1	--	--	2
	%	0.31%	--	1.51%	--	--	0.26%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Neurological - falls (Q13)							
-	N	41	7	2	--	63	113
	%	--	--	--	--	--	--
- Absent	N	172	96	24	25	11	328
	%	365.957	--	--	--	--	697.872
- Recent/active	N	84	35	22	14	25	180
	%	101.205	100.00%	100.00%	100.00%	104.167	101.124
- Remote/inactive	N	17	55	18	14	19	123
	%	106.25	100.00%	105.882	100.00%	100.00%	101.653
Neurological - number of falls (Q13a)		Min	0.00	0.00	0.00	0.00	0.00
		Mean	1.62	0.84	1.67	0.96	1.41
		Std	2.74	0.87	1.88	1.04	2.41
		Max	20.00	4.00	9.00	5.00	25.00
Neurological - other CNS disease (Q14)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	122	139	42	35	37	375
	%	38.85%	72.02%	63.63%	66.03%	31.35%	50.40%
- Recent/active	N	23	32	13	14	14	96
	%	7.32%	16.58%	19.69%	26.41%	11.86%	12.90%
- Remote/inactive	N	6	15	9	4	4	38
	%	1.91%	7.77%	13.63%	7.54%	3.38%	5.10%
- Not determined	N	2	--	--	--	--	2
	%	0.63%	--	--	--	--	0.26%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Common medical - benign prostatic hypertrophy (Q15)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	129	157	40	43	39	408
	%	41.08%	81.34%	60.60%	81.13%	33.05%	54.83%
- Recent/active	N	19	21	18	6	11	75
	%	6.05%	10.88%	27.27%	11.32%	9.32%	10.08%
- Remote/inactive	N	5	7	6	4	5	27
	%	1.59%	3.62%	9.09%	7.54%	4.23%	3.62%
- Not determined	N	--	1	--	--	--	1
	%	--	0.51%	--	--	--	0.13%
Common medical - sleep apnea (Q16)							
	N	129	7	2	--	63	201
	%	41.08%	3.62%	3.03%	--	53.38%	27.01%
- Absent	N	163	151	44	34	39	431
	%	51.91%	78.23%	66.66%	64.15%	33.05%	57.93%
- Recent/active	N	19	31	15	17	13	95
	%	6.05%	16.06%	22.72%	32.07%	11.01%	12.76%
- Remote/inactive	N	2	1	3	--	1	7
	%	0.63%	0.51%	4.54%	--	0.84%	0.94%
- Not determined	N	1	3	2	2	2	10
	%	0.31%	1.55%	3.03%	3.77%	1.69%	1.34%
Common medical - thyroid (Q17)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	100	123	44	36	38	341
	%	31.84%	63.73%	66.66%	67.92%	32.20%	45.83%
- Recent/active	N	49	50	16	15	16	146
	%	15.60%	25.90%	24.24%	28.30%	13.55%	19.62%

(Continued)

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Common medical - thyroid (Q17)							
- Remote/inactive	N	4	13	4	2	--	23
	%	1.27%	6.73%	6.06%	3.77%	--	3.09%
- Not determined	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Common medical - B12 deficiency (Q18)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	137	163	58	44	48	450
	%	43.63%	84.45%	87.87%	83.01%	40.67%	60.48%
- Recent/active	N	12	15	3	6	5	41
	%	3.82%	7.77%	4.54%	11.32%	4.23%	5.51%
- Remote/inactive	N	2	7	2	3	2	16
	%	0.63%	3.62%	3.03%	5.66%	1.69%	2.15%
- Not determined	N	2	1	1	--	--	4
	%	0.63%	0.51%	1.51%	--	--	0.53%
Common medical - diabetes mellitus (Q19)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	136	159	54	46	46	441
	%	43.31%	82.38%	81.81%	86.79%	38.98%	59.27%
- Recent/active	N	15	25	8	6	9	63
	%	4.77%	12.95%	12.12%	11.32%	7.62%	8.46%
- Remote/inactive	N	2	1	1	1	--	5
	%	0.63%	0.51%	1.51%	1.88%	--	0.67%
- Not determined	N	--	1	1	--	--	2
	%	--	0.51%	1.51%	--	--	0.26%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Psychiatric - psychotic disorders (Q20)							
-	N	87	7	2	--	63	159
	%	27.70%	3.62%	3.03%	--	53.38%	21.37%
- Absent	N	222	186	63	50	54	575
	%	70.70%	96.37%	95.45%	94.33%	45.76%	77.28%
- Recent/active	N	4	--	1	2	1	8
	%	1.27%	--	1.51%	3.77%	0.84%	1.07%
- Remote/inactive	N	--	--	--	1	--	1
	%	--	--	--	1.88%	--	0.13%
- Not determined	N	1	--	--	--	--	1
	%	0.31%	--	--	--	--	0.13%
Psychiatric - depression > 2 months (Q21)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	86	142	45	35	31	339
	%	27.38%	73.57%	68.18%	66.03%	26.27%	45.56%
- Recent/active	N	50	22	18	10	13	113
	%	15.92%	11.39%	27.27%	18.86%	11.01%	15.18%
- Remote/inactive	N	16	22	1	8	9	56
	%	5.09%	11.39%	1.51%	15.09%	7.62%	7.52%
- Not determined	N	1	--	--	--	2	3
	%	0.31%	--	--	--	1.69%	0.40%
Psychiatric - anxiety disorders (Q22)							
-	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	236	174	51	46	44	551
	%	75.15%	90.15%	77.27%	86.79%	37.28%	74.05%
- Recent/active	N	12	5	12	7	2	38
	%	3.82%	2.59%	18.18%	13.20%	1.69%	5.10%

(Continued)

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Psychiatric - anxiety disorders (Q22)							
- Remote/inactive	N	8	7	--	--	9	24
	%	2.54%	3.62%	--	--	7.62%	3.22%
- Not determined	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Alcohol / substance - alcohol abuse or dependence (Q23)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	142	178	59	51	51	481
	%	45.22%	92.22%	89.39%	96.22%	43.22%	64.65%
- Recent/active	N	1	1	--	--	1	3
	%	0.31%	0.51%	--	--	0.84%	0.40%
- Remote/inactive	N	10	7	4	2	1	24
	%	3.18%	3.62%	6.06%	3.77%	0.84%	3.22%
- Not determined	N	--	--	1	--	2	3
	%	--	--	1.51%	--	1.69%	0.40%
Alcohol / substance - substance related disorders (non-alcohol) (Q24)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	150	186	61	52	55	504
	%	47.77%	96.37%	92.42%	98.11%	46.61%	67.74%
- Recent/active	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%
- Remote/inactive	N	3	--	1	1	--	5
	%	0.95%	--	1.51%	1.88%	--	0.67%
- Not determined	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Gastrointestinal - GI disease (Q25)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	121	153	51	46	44	415
	%	38.53%	79.27%	77.27%	86.79%	37.28%	55.77%
- Recent/active	N	23	22	7	5	4	61
	%	7.32%	11.39%	10.60%	9.43%	3.38%	8.19%
- Remote/inactive	N	9	11	6	2	7	35
	%	2.86%	5.69%	9.09%	3.77%	5.93%	4.70%
Gastrointestinal - peptic ulcer (Q26)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	237	171	59	49	51	567
	%	75.47%	88.60%	89.39%	92.45%	43.22%	76.20%
- Recent/active	N	4	--	1	--	--	5
	%	1.27%	--	1.51%	--	--	0.67%
- Remote/inactive	N	15	15	4	4	4	42
	%	4.77%	7.77%	6.06%	7.54%	3.38%	5.64%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Sensory - hearing impairment (Q27)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	81	99	24	22	17	243
	%	25.79%	51.29%	36.36%	41.50%	14.40%	32.66%
- Recent/active	N	70	79	37	31	33	250
	%	22.29%	40.93%	56.06%	58.49%	27.96%	33.60%
- Remote/inactive	N	--	2	2	--	1	5
	%	--	1.03%	3.03%	--	0.84%	0.67%
- Not determined	N	2	6	1	--	4	13
	%	0.63%	3.10%	1.51%	--	3.38%	1.74%
Sensory - visual impairment (non-correctable) (Q28)							
-	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	142	180	61	50	50	483
	%	45.22%	93.26%	92.42%	94.33%	42.37%	64.91%
- Recent/active	N	10	6	3	2	3	24
	%	3.18%	3.10%	4.54%	3.77%	2.54%	3.22%
- Remote/inactive	N	--	--	--	1	2	3
	%	--	--	--	1.88%	1.69%	0.40%
- Not determined	N	1	--	--	--	--	1
	%	0.31%	--	--	--	--	0.13%
Sensory - glaucoma (Q29)							
-	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	220	168	53	51	48	540
	%	70.06%	87.04%	80.30%	96.22%	40.67%	72.58%
- Recent/active	N	33	16	8	1	7	65
	%	10.50%	8.29%	12.12%	1.88%	5.93%	8.73%

(Continued)

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Sensory - glaucoma (Q29)							
- Remote/inactive	N	3	1	3	1	--	8
	%	0.95%	0.51%	4.54%	1.88%	--	1.07%
- Not determined	N	--	1	--	--	--	1
	%	--	0.51%	--	--	--	0.13%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Sensory - macular degeneration (Q30)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	236	170	62	47	45	560
	%	75.15%	88.08%	93.93%	88.67%	38.13%	75.26%
- Recent/active	N	18	16	2	6	9	51
	%	5.73%	8.29%	3.03%	11.32%	7.62%	6.85%
- Remote/inactive	N	1	--	--	--	1	2
	%	0.31%	--	--	--	0.84%	0.26%
- Not determined	N	1	--	--	--	--	1
	%	0.31%	--	--	--	--	0.13%
Sensory - cataracts (Q31)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	129	56	18	16	17	236
	%	41.08%	29.01%	27.27%	30.18%	14.40%	31.72%
- Recent/active	N	27	62	12	18	13	132
	%	8.59%	32.12%	18.18%	33.96%	11.01%	17.74%
- Remote/inactive	N	100	67	34	19	25	245
	%	31.84%	34.71%	51.51%	35.84%	21.18%	32.93%
- Not determined	N	--	1	--	--	--	1
	%	--	0.51%	--	--	--	0.13%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Use of aids - wears hearing aids (Q32)							
- .	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- No	N	113	145	44	36	33	371
	%	35.98%	75.12%	66.66%	67.92%	27.96%	49.86%
- Yes	N	40	41	20	17	22	140
	%	12.73%	21.24%	30.30%	32.07%	18.64%	18.81%
Use of aids - wears glasses or corrective lens (Q33)							
- .	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- No	N	23	21	6	10	7	67
	%	7.32%	10.88%	9.09%	18.86%	5.93%	9.00%
- Yes	N	130	165	58	43	48	444
	%	41.40%	85.49%	87.87%	81.13%	40.67%	59.67%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Other medical condition - heart valve disease (Q34)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	228	176	58	48	50	560
	%	72.61%	91.19%	87.87%	90.56%	42.37%	75.26%
- Recent/active	N	11	6	2	2	5	26
	%	3.50%	3.10%	3.03%	3.77%	4.23%	3.49%
- Remote/inactive	N	14	4	3	2	--	23
	%	4.45%	2.07%	4.54%	3.77%	--	3.09%
- Not determined	N	3	--	1	1	--	5
	%	0.95%	--	1.51%	1.88%	--	0.67%
Other medical condition - congestive heart failure (Q35)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	142	185	63	52	53	495
	%	45.22%	95.85%	95.45%	98.11%	44.91%	66.53%
- Recent/active	N	8	--	1	--	1	10
	%	2.54%	--	1.51%	--	0.84%	1.34%
- Remote/inactive	N	3	--	--	--	1	4
	%	0.95%	--	--	--	0.84%	0.53%
- Not determined	N	--	1	--	1	--	2
	%	--	0.51%	--	1.88%	--	0.26%
Other medical condition - other heart conditions (Q36)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	116	137	50	43	37	383
	%	36.94%	70.98%	75.75%	81.13%	31.35%	51.47%
- Recent/active	N	27	32	7	8	13	87
	%	8.59%	16.58%	10.60%	15.09%	11.01%	11.69%

(Continued)

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Other medical condition - other heart conditions (Q36)							
- Remote/inactive	N	10	16	7	--	3	36
	%	3.18%	8.29%	10.60%	--	2.54%	4.83%
- Not determined	N	--	1	--	2	2	5
	%	--	0.51%	--	3.77%	1.69%	0.67%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Other medical condition - asthma, COPD, or other respiratory insufficiency (Q37)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	215	146	54	44	47	506
	%	68.47%	75.64%	81.81%	83.01%	39.83%	68.01%
- Recent/active	N	35	26	9	7	5	82
	%	11.14%	13.47%	13.63%	13.20%	4.23%	11.02%
- Remote/inactive	N	6	14	1	2	3	26
	%	1.91%	7.25%	1.51%	3.77%	2.54%	3.49%
Other medical condition - renal insufficiency (kidney problems) (Q38)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	240	169	57	48	48	562
	%	76.43%	87.56%	86.36%	90.56%	40.67%	75.53%
- Recent/active	N	10	8	3	2	6	29
	%	3.18%	4.14%	4.54%	3.77%	5.08%	3.89%
- Remote/inactive	N	6	9	3	3	1	22
	%	1.91%	4.66%	4.54%	5.66%	0.84%	2.95%
- Not determined	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%
Other medical condition - hepatic insufficiency (liver problems) (Q39)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	253	180	61	51	55	600
	%	80.57%	93.26%	92.42%	96.22%	46.61%	80.64%
- Recent/active	N	2	4	--	2	--	8
	%	0.63%	2.07%	--	3.77%	--	1.07%
- Remote/inactive	N	1	2	1	--	--	4
	%	0.31%	1.03%	1.51%	--	--	0.53%

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
Other medical condition - hepatic insufficiency (liver problems) (Q39)								
- Not determined		N	--	--	2	--	--	2
		%	--	--	3.03%	--	--	0.26%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Other medical condition - osteoarthritis (ordinary arthritis) (Q40)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	109	75	21	14	21	240
	%	34.71%	38.86%	31.81%	26.41%	17.79%	32.25%
- Recent/active	N	145	105	37	36	33	356
	%	46.17%	54.40%	56.06%	67.92%	27.96%	47.84%
- Remote/inactive	N	--	5	5	1	1	12
	%	--	2.59%	7.57%	1.88%	0.84%	1.61%
- Not determined	N	2	1	1	2	--	6
	%	0.63%	0.51%	1.51%	3.77%	--	0.80%
Other medical condition - rheumatoid arthritis / lupus / scleroderma (Q41)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	249	182	60	53	55	599
	%	79.29%	94.30%	90.90%	100.00%	46.61%	80.51%
- Recent/active	N	6	3	2	--	--	11
	%	1.91%	1.55%	3.03%	--	--	1.47%
- Remote/inactive	N	1	1	2	--	--	4
	%	0.31%	0.51%	3.03%	--	--	0.53%
Other medical condition - cancer (Q42)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	168	98	36	29	26	357
	%	53.50%	50.77%	54.54%	54.71%	22.03%	47.98%
- Recent/active	N	20	25	8	5	8	66
	%	6.36%	12.95%	12.12%	9.43%	6.77%	8.87%
- Remote/inactive	N	68	63	18	19	21	189
	%	21.65%	32.64%	27.27%	35.84%	17.79%	25.40%

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
Other medical condition - cancer (Q42)								
- Not determined		N	--	--	2	--	--	2
		%	--	--	3.03%	--	--	0.26%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Other medical condition - STDs (Q43)							
- .	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	253	184	64	51	54	606
	%	80.57%	95.33%	96.96%	96.22%	45.76%	81.45%
- Recent/active	N	--	1	--	--	1	2
	%	--	0.51%	--	--	0.84%	0.26%
- Remote/inactive	N	2	1	--	2	--	5
	%	0.63%	0.51%	--	3.77%	--	0.67%
- Not determined	N	1	--	--	--	--	1
	%	0.31%	--	--	--	--	0.13%
Other medical condition - syphilis (Q43a)							
- .	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	153	186	64	53	55	511
	%	48.72%	96.37%	96.96%	100.00%	46.61%	68.68%
Other medical condition - HIV (Q43b)							
- .	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	153	186	64	53	55	511
	%	48.72%	96.37%	96.96%	100.00%	46.61%	68.68%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Other medical condition - hepatitis B (Q43c)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	148	184	63	52	53	500
	%	47.13%	95.33%	95.45%	98.11%	44.91%	67.20%
- Recent/active	N	1	2	--	--	--	3
	%	0.31%	1.03%	--	--	--	0.40%
- Remote/inactive	N	2	--	--	1	1	4
	%	0.63%	--	--	1.88%	0.84%	0.53%
- Not determined	N	2	--	1	--	1	4
	%	0.63%	--	1.51%	--	0.84%	0.53%
Other medical condition - hepatitis C (Q43d)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	150	186	61	53	54	504
	%	47.77%	96.37%	92.42%	100.00%	45.76%	67.74%
- Remote/inactive	N	1	--	2	--	1	4
	%	0.31%	--	3.03%	--	0.84%	0.53%
- Not determined	N	2	--	1	--	--	3
	%	0.63%	--	1.51%	--	--	0.40%
Other medical condition - other medical (Q44)							
	N	161	7	2	--	63	233
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%
- Absent	N	68	41	16	14	9	148
	%	21.65%	21.24%	24.24%	26.41%	7.62%	19.89%
- Recent/active	N	67	89	34	28	26	244
	%	21.33%	46.11%	51.51%	52.83%	22.03%	32.79%
- Remote/inactive	N	15	56	14	11	19	115
	%	4.77%	29.01%	21.21%	20.75%	16.10%	15.45%
- Not determined	N	3	--	--	--	1	4
	%	0.95%	--	--	--	0.84%	0.53%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Surgeries - open heart surgery / cardiac bypass procedure (Q45)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	232	182	59	52	51	576
	%	73.88%	94.30%	89.39%	98.11%	43.22%	77.41%
- Recent/active	N	1	1	1	--	--	3
	%	0.31%	0.51%	1.51%	--	--	0.40%
- Remote/inactive	N	23	3	4	1	4	35
	%	7.32%	1.55%	6.06%	1.88%	3.38%	4.70%
Surgeries - angioplasty / endarterectomy / stent (Q46)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	238	179	58	48	52	575
	%	75.79%	92.74%	87.87%	90.56%	44.06%	77.28%
- Recent/active	N	4	1	2	2	1	10
	%	1.27%	0.51%	3.03%	3.77%	0.84%	1.34%
- Remote/inactive	N	14	6	4	3	2	29
	%	4.45%	3.10%	6.06%	5.66%	1.69%	3.89%
Surgeries - hysterectomy / oophorectomy (Q47)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	199	136	48	45	42	470
	%	63.37%	70.46%	72.72%	84.90%	35.59%	63.17%
- Recent/active	N	1	3	3	1	1	9
	%	0.31%	1.55%	4.54%	1.88%	0.84%	1.20%
- Remote/inactive	N	56	47	13	7	12	135
	%	17.83%	24.35%	19.69%	13.20%	10.16%	18.14%
Surgeries - other surgeries (Q48)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- Absent	N	89	13	2	4	3	111

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Surgeries - other surgeries (Q48)							
- Absent	%	28.34%	6.73%	3.03%	7.54%	2.54%	14.91%
- Recent/active	N	34	39	18	12	17	120
	%	10.82%	20.20%	27.27%	22.64%	14.40%	16.12%
- Remote/inactive	N	132	134	44	37	35	382
	%	42.03%	69.43%	66.66%	69.81%	29.66%	51.34%
- Not determined	N	1	--	--	--	--	1
	%	0.31%	--	--	--	--	0.13%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Hospitalizations (Q49)							
	N	58	7	2	--	63	130
	%	18.47%	3.62%	3.03%	--	53.38%	17.47%
- No	N	190	157	49	44	44	484
	%	60.50%	81.34%	74.24%	83.01%	37.28%	65.05%
- Yes	N	66	29	15	9	11	130
	%	21.01%	15.02%	22.72%	16.98%	9.32%	17.47%

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
		314	193	66	53	118	744	
Smoking history - subject has ever smoked (Q50)								
- .	N	161	7	2	--	63	233	
	%	51.27%	3.62%	3.03%	--	53.38%	31.31%	
- No	N	84	111	30	29	31	285	
	%	26.75%	57.51%	45.45%	54.71%	26.27%	38.30%	
- Yes	N	69	75	34	24	24	226	
	%	21.97%	38.86%	51.51%	45.28%	20.33%	30.37%	
Smoking history - smoked in the last 30 days (Q50a)								
- .	N	245	118	32	29	94	518	
	%	78.02%	61.13%	48.48%	54.71%	79.66%	69.62%	
- No	N	63	75	33	24	24	219	
	%	20.06%	38.86%	50.00%	45.28%	20.33%	29.43%	
- Yes	N	6	--	1	--	--	7	
	%	1.91%	--	1.51%	--	--	0.94%	
Smoking history - quit smoking (Q50b)								
- .	N	245	118	32	29	94	518	
	%	78.02%	61.13%	48.48%	54.71%	79.66%	69.62%	
- No	N	5	1	1	--	--	7	
	%	1.59%	0.51%	1.51%	--	--	0.94%	
- Yes	N	64	74	33	24	24	219	
	%	20.38%	38.34%	50.00%	45.28%	20.33%	29.43%	
Smoking history - age quit smoking (Q50b1)		Min	19.00	20.00	20.00	11.00	21.00	11.00
	Mean	43.83	35.42	36.88	38.71	37.71	38.71	
	Std	12.85	10.20	11.58	15.68	13.49	12.61	
	Max	74.00	66.00	66.00	68.00	70.00	74.00	
Smoking history - total years smoked (Q50c)		Min	1.00	1.00	1.00	1.00	1.00	
	Mean	19.95	13.49	14.28	20.29	15.75	16.56	
	Std	12.34	9.15	11.48	16.80	12.21	12.09	
	Max	47.00	41.00	40.00	52.00	50.00	52.00	
Smoking history - average number of packs per day (Q50d)								
- .	N	245	118	32	29	94	518	

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Smoking history - average number of packs per day (Q50d)							
- .	%	78.02%	61.13%	48.48%	54.71%	79.66%	69.62%
- one cigarette to less than half pack	N	32	24	14	13	9	92
	%	10.19%	12.43%	21.21%	24.52%	7.62%	12.36%
- half pack to one and half pack	N	22	32	12	6	7	79
	%	7.00%	16.58%	18.18%	11.32%	5.93%	10.61%
- one and half pack to two packs	N	5	7	4	3	3	22
	%	1.59%	3.62%	6.06%	5.66%	2.54%	2.95%
- more than two packs	N	5	4	2	--	--	11
	%	1.59%	2.07%	3.03%	--	--	1.47%
- Unable to remember / not determined	N	5	8	2	2	5	22
	%	1.59%	4.14%	3.03%	3.77%	4.23%	2.95%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Alcohol use (past year) - subject currently uses alcohol (Q51)							
	N	129	7	2	--	63	201
	%	41.08%	3.62%	3.03%	--	53.38%	27.01%
- No							
	N	91	53	18	19	18	199
	%	28.98%	27.46%	27.27%	35.84%	15.25%	26.74%
- Yes							
	N	94	133	46	34	37	344
	%	29.93%	68.91%	69.69%	64.15%	31.35%	46.23%
Alcohol use (past year) - average # drinks per day: hard liquor (Q51a1)		Min	0.00	0.00	0.00	0.00	0.00
	Mean	0.10	0.06	0.03	0.07	0.06	0.07
	Std	0.36	0.32	0.16	0.26	0.34	0.31
	Max	2.00	2.00	1.00	1.00	2.00	2.00
Alcohol use (past year) - average # drinks per day: wine (Q51a2)		Min	0.00	0.00	0.00	0.00	0.00
	Mean	0.48	0.48	0.41	0.33	0.57	0.47
	Std	0.80	0.73	0.59	0.55	0.78	0.72
	Max	4.00	3.00	2.00	2.00	3.00	4.00
Alcohol use (past year) - average # drinks per day: beer (Q51a3)		Min	0.00	0.00	0.00	0.00	0.00
	Mean	0.03	0.03	0.03	0.04	0.06	0.03
	Std	0.31	0.17	0.16	0.19	0.24	0.23
	Max	3.00	1.00	1.00	1.00	1.00	3.00

Medical History		Status					All	
		.	Normal	MCI	QCI	Dementia		
		314	193	66	53	118	744	
Alcohol use (past year) - greater than 4 or 5 drinks on occasion (Q51b)								
	N	220	60	20	19	80	399	
	%	70.06%	31.08%	30.30%	35.84%	67.79%	53.62%	
	- No	N	88	120	45	30	36	319
		%	28.02%	62.17%	68.18%	56.60%	30.50%	42.87%
	- Yes	N	6	13	--	4	1	24
		%	1.91%	6.73%	--	7.54%	0.84%	3.22%
	- Not determined	N	--	--	1	--	1	2
		%	--	--	1.51%	--	0.84%	0.26%
Alcohol use (past year) - increased cognitive difficulties (Q51c1)								
	N	220	60	20	19	80	399	
	%	70.06%	31.08%	30.30%	35.84%	67.79%	53.62%	
	- No	N	85	130	44	31	37	327
		%	27.07%	67.35%	66.66%	58.49%	31.35%	43.95%
	- Yes	N	9	1	2	3	1	16
		%	2.86%	0.51%	3.03%	5.66%	0.84%	2.15%
	- Not determined	N	--	2	--	--	--	2
		%	--	1.03%	--	--	--	0.26%
Alcohol use (past year) - personality changes (Q51c2)								
	N	220	60	20	19	80	399	
	%	70.06%	31.08%	30.30%	35.84%	67.79%	53.62%	
	- No	N	87	130	45	33	35	330
		%	27.70%	67.35%	68.18%	62.26%	29.66%	44.35%
	- Yes	N	7	1	1	1	3	13
		%	2.22%	0.51%	1.51%	1.88%	2.54%	1.74%
	- Not determined	N	--	2	--	--	--	2
		%	--	1.03%	--	--	--	0.26%
Alcohol use (past year) - loss of balance (Q51c3)								
	N	220	60	20	19	80	399	
	%	70.06%	31.08%	30.30%	35.84%	67.79%	53.62%	

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Alcohol use (past year) - loss of balance (Q51c3)							
- No	N	90	129	44	34	36	333
	%	28.66%	66.83%	66.66%	64.15%	30.50%	44.75%
- Yes	N	4	2	2	--	2	10
	%	1.27%	1.03%	3.03%	--	1.69%	1.34%
- Not determined	N	--	2	--	--	--	2
	%	--	1.03%	--	--	--	0.26%

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
		314	193	66	53	118	744
Past alcohol use - subject used alcohol in the previous 5 years (Q52)							
- .	N	129	7	2	--	63	201
	%	41.08%	3.62%	3.03%	--	53.38%	27.01%
- No	N	79	44	14	14	12	163
	%	25.15%	22.79%	21.21%	26.41%	10.16%	21.90%
- Yes	N	106	142	50	39	43	380
	%	33.75%	73.57%	75.75%	73.58%	36.44%	51.07%
Past alcohol use - how much drink on average (Q52a)							
- .	N	208	51	16	14	75	364
	%	66.24%	26.42%	24.24%	26.41%	63.55%	48.92%
- 0-7 drinks per week	N	90	121	42	36	38	327
	%	28.66%	62.69%	63.63%	67.92%	32.20%	43.95%
- 8-14 drinks per week	N	12	17	7	1	4	41
	%	3.82%	8.80%	10.60%	1.88%	3.38%	5.51%
- 15-21 drinks per week	N	4	3	--	2	1	10
	%	1.27%	1.55%	--	3.77%	0.84%	1.34%
- 22-28 drinks per week	N	--	1	--	--	--	1
	%	--	0.51%	--	--	--	0.13%
- > 30 drinks per week	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%
Past alcohol use - stopped drinking (Q52b)							
- .	N	208	51	16	14	75	364
	%	66.24%	26.42%	24.24%	26.41%	63.55%	48.92%
- No	N	89	136	46	34	37	342
	%	28.34%	70.46%	69.69%	64.15%	31.35%	45.96%
- Yes	N	17	6	4	5	6	38
	%	5.41%	3.10%	6.06%	9.43%	5.08%	5.10%
Past alcohol abuse - subject has a history of alcohol abuse (Q53)							
- .	N	147	7	2	--	63	219
	%	46.81%	3.62%	3.03%	--	53.38%	29.43%

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in in summary.

Medical History		Status					All
		.	Normal	MCI	QCI	Dementia	
Past alcohol abuse - subject has a history of alcohol abuse (Q53)							
- No	N	155	178	59	50	53	495
	%	49.36%	92.22%	89.39%	94.33%	44.91%	66.53%
- Yes	N	12	8	4	3	2	29
	%	3.82%	4.14%	6.06%	5.66%	1.69%	3.89%
- Not determined	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%