

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Subject reports a decline in memory (Q1)							
-	N	3	--	--	--	--	3
	%	14.28%	--	--	--	--	0.66%
- No	N	1	166	15	20	23	225
	%	4.76%	85.56%	22.72%	37.73%	19.49%	49.77%
- Yes	N	7	28	51	33	50	169
	%	33.33%	14.43%	77.27%	62.26%	42.37%	37.38%
- 8	N	10	--	--	--	45	55
	%	47.61%	--	--	--	38.13%	12.16%
Information report a decline in subject's memory (Q2)							
-	N	3	--	--	--	--	3
	%	14.28%	--	--	--	--	0.66%
- No	N	--	173	11	27	2	213
	%	--	89.17%	16.66%	50.94%	1.69%	47.12%
- Yes	N	18	21	55	26	116	236
	%	85.71%	10.82%	83.33%	49.05%	98.30%	52.21%
At what age cognitive declined (Q3b)							
	Min	50.00	66.00	52.00	54.00	47.00	47.00
	Mean	63.44	74.25	72.88	71.52	70.73	71.03
	Std	8.51	5.79	7.24	8.97	9.47	8.90
	Max	83.00	86.00	88.00	92.00	95.00	95.00
Cognitive symptoms of memory (Q4a)							
-	N	3	182	--	28	--	213
	%	14.28%	93.81%	--	52.83%	--	47.12%
- No	N	--	9	7	7	--	23
	%	--	4.63%	10.60%	13.20%	--	5.08%
- Yes	N	18	3	59	18	118	216
	%	85.71%	1.54%	89.39%	33.96%	100.00%	47.78%
Cognitive symptoms of judgment/problem solving (Q4b)							
-	N	3	182	--	28	--	213
	%	14.28%	93.81%	--	52.83%	--	47.12%

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
Cognitive symptoms of judgment/problem solving (Q4b)							
- No	N	--	12	38	21	4	75
	%	--	6.18%	57.57%	39.62%	3.38%	16.59%
- Yes	N	18	--	28	4	114	164
	%	85.71%	--	42.42%	7.54%	96.61%	36.28%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Clinician Judgment of Symptoms		Status				
		.	Normal	MCI	QCI	Dementia
		21	194	66	53	118
Cognitive symptoms of language (Q4c)						
- .	N	3	182	--	28	--
	%	14.28%	93.81%	--	52.83%	--
- No	N	--	2	24	15	4
	%	--	1.03%	36.36%	28.30%	3.38%
- Yes	N	18	10	42	10	114
	%	85.71%	5.15%	63.63%	18.86%	96.61%
Cognitive symptoms of visuospatial function (Q4d)						
- .	N	3	182	--	28	--
	%	14.28%	93.81%	--	52.83%	--
- No	N	4	12	54	24	34
	%	19.04%	6.18%	81.81%	45.28%	28.81%
- Yes	N	14	--	12	1	84
	%	66.66%	--	18.18%	1.88%	71.18%
Cognitive symptoms of attention/concentration (Q4e)						
- .	N	3	182	--	28	--
	%	14.28%	93.81%	--	52.83%	--
- No	N	2	9	39	20	19
	%	9.52%	4.63%	59.09%	37.73%	16.10%
- Yes	N	16	3	27	5	99
	%	76.19%	1.54%	40.90%	9.43%	83.89%
Cognitive symptoms of fluctuating cognition (Q4f)						
- .	N	3	182	--	28	--
	%	14.28%	93.81%	--	52.83%	--
- No	N	9	12	66	25	109
	%	42.85%	6.18%	100.00%	47.16%	92.37%
- Yes	N	7	--	--	--	8
	%	33.33%	--	--	--	6.77%
- Unknown	N	2	--	--	--	1
	%	9.52%	--	--	--	0.84%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Clinician Judgment of Symptoms		All
		452
Cognitive symptoms of language (Q4c)		
- .	N	213
	%	47.12%
- No	N	45
	%	9.95%
- Yes	N	194
	%	42.92%
Cognitive symptoms of visuospatial function (Q4d)		
- .	N	213
	%	47.12%
- No	N	128
	%	28.31%
- Yes	N	111
	%	24.55%
Cognitive symptoms of attention/concentration (Q4e)		
- .	N	213
	%	47.12%
- No	N	89
	%	19.69%
- Yes	N	150
	%	33.18%
Cognitive symptoms of fluctuating cognition (Q4f)		
- .	N	213
	%	47.12%
- No	N	221
	%	48.89%
- Yes	N	15
	%	3.31%
- Unknown	N	3
	%	0.66%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Mode of onset of cognitive symptoms (Q6)							
	N	3	182	--	28	--	213
	%	14.28%	93.81%	--	52.83%	--	47.12%
- Gradual (> 6 months)	N	18	9	64	22	118	231
	%	85.71%	4.63%	96.96%	41.50%	100.00%	51.10%
- Unknown	N	--	3	2	3	--	8
	%	--	1.54%	3.03%	5.66%	--	1.76%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Mode of onset of cognitive symptoms (Q6)							
	N	3	182	--	28	--	213
	%	14.28%	93.81%	--	52.83%	--	47.12%
- Gradual (> 6 months)	N	18	9	64	22	118	231
	%	85.71%	4.63%	96.96%	41.50%	100.00%	51.10%
- Unknown	N	--	3	2	3	--	8
	%	--	1.54%	3.03%	5.66%	--	1.76%

Clinician Judgment of Symptoms		Status				
		.	Normal	MCI	QCI	Dementia
		21	194	66	53	118
Behavior symptoms of apathy (Q7a)						
	N	5	160	23	28	22
	%	23.80%	82.47%	34.84%	52.83%	18.64%
- No	N	6	27	32	16	36
	%	28.57%	13.91%	48.48%	30.18%	30.50%
- Yes	N	9	7	11	9	56
	%	42.85%	3.60%	16.66%	16.98%	47.45%
- Unknown	N	1	--	--	--	4
	%	4.76%	--	--	--	3.38%
Behavior symptoms of depression (HPC Q7b)						
	N	5	160	23	28	22
	%	23.80%	82.47%	34.84%	52.83%	18.64%
- No	N	5	16	20	13	59
	%	23.80%	8.24%	30.30%	24.52%	50.00%
- Yes	N	10	18	23	12	34
	%	47.61%	9.27%	34.84%	22.64%	28.81%
- Unknown	N	1	--	--	--	3
	%	4.76%	--	--	--	2.54%
Behavior symptoms of visual hallucination (Q7c1)						
	N	5	160	23	28	22
	%	23.80%	82.47%	34.84%	52.83%	18.64%
- No	N	10	34	43	25	76
	%	47.61%	17.52%	65.15%	47.16%	64.40%
- Yes	N	6	--	--	--	15
	%	28.57%	--	--	--	12.71%
- Unknown	N	--	--	--	--	5
	%	--	--	--	--	4.23%
Behavior symptoms of auditory hallucination (Q7c2)						
	N	5	160	23	28	22
	%	23.80%	82.47%	34.84%	52.83%	18.64%
- No	N	13	34	43	24	79

(Continued)

Clinician Judgment of Symptoms		All
		452
Behavior symptoms of apathy (Q7a)		
- .	N	238
	%	52.65%
- No	N	117
	%	25.88%
- Yes	N	92
	%	20.35%
- Unknown	N	5
	%	1.10%
Behavior symptoms of depression (HPC Q7b)		
- .	N	238
	%	52.65%
- No	N	113
	%	25.00%
- Yes	N	97
	%	21.46%
- Unknown	N	4
	%	0.88%
Behavior symptoms of visual hallucination (Q7c1)		
- .	N	238
	%	52.65%
- No	N	188
	%	41.59%
- Yes	N	21
	%	4.64%
- Unknown	N	5
	%	1.10%
Behavior symptoms of auditory hallucination (Q7c2)		
- .	N	238
	%	52.65%
- No	N	193

(Continued)

Clinician Judgment of Symptoms		Status				
		.	Normal	MCI	QCI	Dementia
Behavior symptoms of auditory hallucination (Q7c2)						
- No	%	61.90%	17.52%	65.15%	45.28%	66.94%
- Yes	N	2	--	--	1	9
	%	9.52%	--	--	1.88%	7.62%
- Unknown	N	1	--	--	--	8
	%	4.76%	--	--	--	6.77%

Clinician Judgment of Symptoms		All
Behavior symptoms of auditory hallucination (Q7c2)		452
- No	%	42.69%
- Yes	N	12
	%	2.65%
- Unknown	N	9
	%	1.99%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Behavior symptoms of abnormal/false/delusional beliefs (Q7c3)							
	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- No	N	11	32	43	24	76	186
	%	52.38%	16.49%	65.15%	45.28%	64.40%	41.15%
- Yes	N	4	2	--	1	14	21
	%	19.04%	1.03%	--	1.88%	11.86%	4.64%
- Unknown	N	1	--	--	--	6	7
	%	4.76%	--	--	--	5.08%	1.54%
Behavior symptoms of disinhibition (Q7d)							
	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- No	N	13	32	39	25	63	172
	%	61.90%	16.49%	59.09%	47.16%	53.38%	38.05%
- Yes	N	3	2	4	--	31	40
	%	14.28%	1.03%	6.06%	--	26.27%	8.84%
- Unknown	N	--	--	--	--	2	2
	%	--	--	--	--	1.69%	0.44%
Behavior symptoms of irritability (Q7e)							
	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- No	N	7	27	19	15	60	128
	%	33.33%	13.91%	28.78%	28.30%	50.84%	28.31%
- Yes	N	8	7	24	10	35	84
	%	38.09%	3.60%	36.36%	18.86%	29.66%	18.58%
- Unknown	N	1	--	--	--	1	2
	%	4.76%	--	--	--	0.84%	0.44%
Behavior symptoms of agitation (Q7f)							
	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- No	N	5	34	31	22	56	148

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
Behavior symptoms of agitation (Q7f)							
- No	%	23.80%	17.52%	46.96%	41.50%	47.45%	32.74%
- Yes	N	11	--	12	3	39	65
	%	52.38%	--	18.18%	5.66%	33.05%	14.38%
- Unknown	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.22%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Behavior symptoms of personality change (Q7g)							
-	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- No	N	10	30	39	20	67	166
	%	47.61%	15.46%	59.09%	37.73%	56.77%	36.72%
- Yes	N	6	4	4	5	29	48
	%	28.57%	2.06%	6.06%	9.43%	24.57%	10.61%
Behavior symptoms of REM sleep behavior disorder (Q7h)							
-	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- No	N	12	32	41	24	92	201
	%	57.14%	16.49%	62.12%	45.28%	77.96%	44.46%
- Yes	N	4	2	2	1	2	11
	%	19.04%	1.03%	3.03%	1.88%	1.69%	2.43%
- Unknown	N	--	--	--	--	2	2
	%	--	--	--	--	1.69%	0.44%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Mode of onset of behavioral symptoms (Q9)							
	N	5	160	23	28	22	238
	%	23.80%	82.47%	34.84%	52.83%	18.64%	52.65%
- Gradual (> 6 months)	N	15	23	41	22	90	191
	%	71.42%	11.85%	62.12%	41.50%	76.27%	42.25%
- Subacute (= < 6 months)	N	1	1	--	1	3	6
	%	4.76%	0.51%	--	1.88%	2.54%	1.32%
- Abrupt (within days)	N	--	4	--	--	2	6
	%	--	2.06%	--	--	1.69%	1.32%
- Other	N	--	1	--	--	--	1
	%	--	0.51%	--	--	--	0.22%
- Unknown	N	--	5	2	2	1	10
	%	--	2.57%	3.03%	3.77%	0.84%	2.21%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Motor symptoms of gait disorder (Q10a)							
	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- No	N	1	35	19	9	25	89
	%	4.76%	18.04%	28.78%	16.98%	21.18%	19.69%
- Yes	N	14	27	27	15	72	155
	%	66.66%	13.91%	40.90%	28.30%	61.01%	34.29%
Motor symptoms of falls (Q10b)							
	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- No	N	7	55	40	24	71	197
	%	33.33%	28.35%	60.60%	45.28%	60.16%	43.58%
- Yes	N	8	7	6	--	25	46
	%	38.09%	3.60%	9.09%	--	21.18%	10.17%
- Unknown	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.22%
Motor symptoms of tremor (Q10c)							
	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- No	N	8	22	13	8	34	85
	%	38.09%	11.34%	19.69%	15.09%	28.81%	18.80%
- Yes	N	7	40	33	16	62	158
	%	33.33%	20.61%	50.00%	30.18%	52.54%	34.95%
- Unknown	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.22%
Motor symptoms of slowness (Q10d)							
	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- No	N	--	46	25	16	26	113
	%	--	23.71%	37.87%	30.18%	22.03%	25.00%

(Continued)

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
Motor symptoms of slowness (Q10d) - Yes	N	15	16	21	8	71	131
	%	71.42%	8.24%	31.81%	15.09%	60.16%	28.98%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
First recognized symptom as a decline in motor symptoms (Q11)							
-	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- Gait disorder	N	8	19	15	10	31	83
	%	38.09%	9.79%	22.72%	18.86%	26.27%	18.36%
- Falls	N	--	1	1	--	5	7
	%	--	0.51%	1.51%	--	4.23%	1.54%
- Tremor	N	2	32	20	11	32	97
	%	9.52%	16.49%	30.30%	20.75%	27.11%	21.46%
- Slowness	N	5	8	9	3	25	50
	%	23.80%	4.12%	13.63%	5.66%	21.18%	11.06%
- Unknown	N	--	2	1	--	4	7
	%	--	1.03%	1.51%	--	3.38%	1.54%
Mode of onset of motor symptoms (Q12)							
-	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- Gradual (> 6 months)	N	15	55	45	20	86	221
	%	71.42%	28.35%	68.18%	37.73%	72.88%	48.89%
- Subacute (=< 6 months)	N	--	--	--	2	4	6
	%	--	--	--	3.77%	3.38%	1.32%
- Abrupt (within days)	N	--	1	--	--	2	3
	%	--	0.51%	--	--	1.69%	0.66%
- Other	N	--	3	--	2	1	6
	%	--	1.54%	--	3.77%	0.84%	1.32%
- Unknown	N	--	3	1	--	4	8
	%	--	1.54%	1.51%	--	3.38%	1.76%

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
		21	194	66	53	118	452
Suggestive of parkinsonism (Q12a)							
-	N	6	132	20	29	21	208
	%	28.57%	68.04%	30.30%	54.71%	17.79%	46.01%
- No	N	4	59	35	23	59	180
	%	19.04%	30.41%	53.03%	43.39%	50.00%	39.82%
- Yes	N	10	2	7	1	25	45
	%	47.61%	1.03%	10.60%	1.88%	21.18%	9.95%
- 9	N	1	1	4	--	13	19
	%	4.76%	0.51%	6.06%	--	11.01%	4.20%
course of overall cognitive/behavioral/motor syndrome (Q13)							
-	N	3	--	--	--	--	3
	%	14.28%	--	--	--	--	0.66%
- Gradually	N	17	70	62	41	117	307
	%	80.95%	36.08%	93.93%	77.35%	99.15%	67.92%
- Stepwise	N	--	2	--	--	--	2
	%	--	1.03%	--	--	--	0.44%
- Static	N	--	3	3	1	--	7
	%	--	1.54%	4.54%	1.88%	--	1.54%
- Fluctuating	N	1	--	--	1	1	3
	%	4.76%	--	--	1.88%	0.84%	0.66%
- Improved	N	--	1	--	--	--	1
	%	--	0.51%	--	--	--	0.22%
- 8	N	--	106	--	7	--	113
	%	--	54.63%	--	13.20%	--	25.00%
- Unknown	N	--	12	1	3	--	16
	%	--	6.18%	1.51%	5.66%	--	3.53%
First recognized change in subject (Q14)							
-	N	3	--	--	--	--	3
	%	14.28%	--	--	--	--	0.66%
- Cognition	N	15	8	49	18	109	199
	%	71.42%	4.12%	74.24%	33.96%	92.37%	44.02%

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Clinician Judgment of Symptoms		Status					All
		.	Normal	MCI	QCI	Dementia	
First recognized change in subject (Q14)							
- Behavior	N	1	27	8	15	6	57
	%	4.76%	13.91%	12.12%	28.30%	5.08%	12.61%
- Motor function	N	2	53	9	13	3	80
	%	9.52%	27.31%	13.63%	24.52%	2.54%	17.69%
- 8	N	--	106	--	7	--	113
	%	--	54.63%	--	13.20%	--	25.00%