Mach: lith.			Sta	atus		All
Mobility		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Independence when walking (MBLTY Q1)						
Name	N	178	53	41	38	310
- Normal	%	92.70%	73.61%	77.35%	42.22%	76.16%
Walles aloud out a support	N	7	12	9	26	54
- Walks slowly w/o support		3.64%	16.66%	16.98%	28.88%	13.26%
	N	7	7	3	23	40
- Frequently uses supportive device	%	3.64%	9.72%	5.66%	25.55%	9.82%
Description who wind a company from a the co	N				3	3
- Requires physical support from others	%				3.33%	0.73%
Endurance when walking (MBLTY Q2)						
- Normal	N	167	55	44	45	311
- Normai	%	86.97%	76.38%	83.01%	50.00%	76.41%
Only able to wall about distances	N	23	13	6	35	77
- Only able to walk short distances	%	11.97%	18.05%	11.32%	38.88%	18.91%
Ciana of and wones when welling about distance	N	2	4	2	9	17
- Signs of endurence when walking short distances		1.04%	5.55%	3.77%	10.00%	4.17%
Linchia to walls make their a face at a second	N			1	1	2
- Unable to walk more than a few steps	%			1.88%	1.11%	0.49%

Mobility	Mobility		Sta	atus		All
Mobility	-	Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Use of a cane (MBLTY Q3a)						
Ne	N	180	60	48	73	361
- No	%	93.75%	83.33%	90.56%	81.11%	88.69%
V	N	12	12	5	17	46
- Yes	%	6.25%	16.66%	9.43%	18.88%	11.30%

Mobility			Sta	atus		All
Widding		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Type of cane use (MBLTY Q3a)						
	N	180	60	48	73	361
	%	93.75%	83.33%	90.56%	81.11%	88.69%
Chronia	N	5	5	1	10	21
- Chronic	%	2.60%	6.94%	1.88%	11.11%	5.15%
		7	7	4	7	25
- Acute	%	3.64%	9.72%	7.54%	7.77%	6.14%

Mobility	Mobility		Sta	atus		All
Mobility		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Use of a walker (MBLTY Q3b)						
- No	N	184	66	50	71	371
-110	%	95.83%	91.66%	94.33%	78.88%	91.15%
V	N	8	6	3	19	36
- Yes	%	4.16%	8.33%	5.66%	21.11%	8.84%

Mobility			All			
		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Type of walker use (MBLTY Q3b)						
	N	184	66	50	71	371
	%	95.83%	91.66%	94.33%	78.88%	91.15%

(Continued)

Mobility				All		
		Normal	MCI	QCI	Dementia	All
Type of walker use (MBLTY Q3b) - Chronic	N	2	3	3	14	22
	%	1.04%	4.16%	5.66%	15.55%	5.40%
	N	6	3		5	14
- Acute	%	3.12%	4.16%		5.55%	3.43%

Mobility				All		
Mobility	y		MCI	QCI	Dementia	All
		192	72	53	90	407
Use of a wheelchair (MBLTY Q3c)						
No	N	192	70	52	85	399
- No	%	100.00%	97.22%	98.11%	94.44%	98.03%
- Yes	N		2	1	5	8
	%		2.77%	1.88%	5.55%	1.96%

Mobility			Sta	itus		All
Mobility		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Type of wheel chair use (MBLTY Q3c)						
	N	192	70	52	85	399
- .	%	100.00%	97.22%	98.11%	94.44%	98.03%
Chronic	N		1		2	3
- Chronic	%		1.38%		2.22%	0.73%
			1	1	3	5
- Acute	%		1.38%	1.88%	3.33%	1.22%

Mahilih	Mobility		Sta	atus		All
Mobility		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Use of bath rails (MBLTY Q3d)						
- No	N	175	61	48	59	343
- 100	%	91.14%	84.72%	90.56%	65.55%	84.27%
V	N	17	11	5	31	64
- Yes	%	8.85%	15.27%	9.43%	34.44%	15.72%

Mobility				All		
		Normal	MCI	QCI	Dementia	All
		192	72	53	90	407
Type of bath rails use (MBLTY Q3d)						
	N	175	61	48	59	343
	%	91.14%	84.72%	90.56%	65.55%	84.27%

(Continued)

Mobility				All		
		Normal	MCI	QCI	Dementia	All
Type of bath rails use (MBLTY Q3d) - Chronic	N	11	9	5	23	48
	%	5.72%	12.50%	9.43%	25.55%	11.79%
	N	6	2		8	16
- Acute		3.12%	2.77%		8.88%	3.93%

Clinic 3.0 Data: 19 January 2021

Mobility				All		
Mobility			MCI	QCI	Dementia	All
		192	72	53	90	407
Difficulty climbing stairs (MBLTY Q4)						
No	N	183	64	47	62	356
- No	%	95.31%	88.88%	88.67%	68.88%	87.46%
W ₂ -	N	9	8	6	28	51
- Yes		4.68%	11.11%	11.32%	31.11%	12.53%