

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject has ever taken Aricept (MTH Q1)							
-	N	3	2	--	--	8	13
	%	0.90%	1.03%	--	--	6.77%	1.70%
- No	N	165	189	42	47	26	469
	%	49.69%	97.92%	63.63%	88.67%	22.03%	61.54%
- Yes	N	164	2	24	6	84	280
	%	49.39%	1.03%	36.36%	11.32%	71.18%	36.74%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject still taking Aricept (MTH Q1A)							
-	N	168	191	42	47	34	482
	%	50.60%	98.96%	63.63%	88.67%	28.81%	63.25%
- No	N	88	2	10	3	36	139
	%	26.50%	1.03%	15.15%	5.66%	30.50%	18.24%
- Yes	N	76	--	14	3	48	141
	%	22.89%	--	21.21%	5.66%	40.67%	18.50%

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
		332	193	66	53	118
Reason for stopping Aricept - Did not help						
- .	N	244	191	56	50	82
	%	73.49%	98.96%	84.84%	94.33%	69.49%
- No	N	52	2	8	1	25
	%	15.66%	1.03%	12.12%	1.88%	21.18%
- Yes	N	36	--	2	2	11
	%	10.84%	--	3.03%	3.77%	9.32%
Reason for stopping Aricept - Abdominal Pain/Cramps						
- .	N	244	191	56	50	82
	%	73.49%	98.96%	84.84%	94.33%	69.49%
- No	N	84	2	9	3	31
	%	25.30%	1.03%	13.63%	5.66%	26.27%
- Yes	N	4	--	1	--	5
	%	1.20%	--	1.51%	--	4.23%
Reason for stopping Aricept - Agitation/Restlessness						
- .	N	244	191	56	50	82
	%	73.49%	98.96%	84.84%	94.33%	69.49%
- No	N	85	2	7	3	31
	%	25.60%	1.03%	10.60%	5.66%	26.27%
- Yes	N	3	--	3	--	5
	%	0.90%	--	4.54%	--	4.23%
Reason for stopping Aricept - Anorexia						
- .	N	244	191	56	50	82
	%	73.49%	98.96%	84.84%	94.33%	69.49%
- No	N	88	2	10	3	36
	%	26.50%	1.03%	15.15%	5.66%	30.50%
Reason for stopping Aricept - Confusion						
- .	N	244	191	56	50	82
	%	73.49%	98.96%	84.84%	94.33%	69.49%
- No	N	85	2	8	3	34
	%	25.60%	1.03%	12.12%	5.66%	28.81%

(Continued)

Medication Treatment History		All
		762
Reason for stopping Aricept - Did not help		
- .	N	623
	%	81.75%
- No	N	88
	%	11.54%
- Yes	N	51
	%	6.69%
Reason for stopping Aricept - Abdominal Pain/Cramps		
- .	N	623
	%	81.75%
- No	N	129
	%	16.92%
- Yes	N	10
	%	1.31%
Reason for stopping Aricept - Agitation/Restlessness		
- .	N	623
	%	81.75%
- No	N	128
	%	16.79%
- Yes	N	11
	%	1.44%
Reason for stopping Aricept - Anorexia		
- .	N	623
	%	81.75%
- No	N	139
	%	18.24%
Reason for stopping Aricept - Confusion		
- .	N	623
	%	81.75%
- No	N	132
	%	17.32%

(Continued)

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
Reason for stopping Aricept - Confusion						
- Yes	N	3	--	2	--	2
	%	0.90%	--	3.03%	--	1.69%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		All
Reason for stopping Aricept - Confusion		762
- Yes	N	7
	%	0.91%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Aricept - Diarrhea							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	78	1	10	3	33	125
	%	23.49%	0.51%	15.15%	5.66%	27.96%	16.40%
- Yes	N	10	1	--	--	3	14
	%	3.01%	0.51%	--	--	2.54%	1.83%
Reason for stopping Aricept - Dizziness							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	82	2	8	3	32	127
	%	24.69%	1.03%	12.12%	5.66%	27.11%	16.66%
- Yes	N	6	--	2	--	4	12
	%	1.80%	--	3.03%	--	3.38%	1.57%
Reason for stopping Aricept - Fatigue							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	88	2	8	3	35	136
	%	26.50%	1.03%	12.12%	5.66%	29.66%	17.84%
- Yes	N	--	--	2	--	1	3
	%	--	--	3.03%	--	0.84%	0.39%
Reason for stopping Aricept - Headache							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	87	2	9	3	35	136
	%	26.20%	1.03%	13.63%	5.66%	29.66%	17.84%
- Yes	N	1	--	1	--	1	3
	%	0.30%	--	1.51%	--	0.84%	0.39%
Reason for stopping Aricept - Insomnia							

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Aricept - Insomnia							
-	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	87	2	9	3	35	136
	%	26.20%	1.03%	13.63%	5.66%	29.66%	17.84%
- Yes	N	1	--	1	--	1	3
	%	0.30%	--	1.51%	--	0.84%	0.39%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Aricept - Nightmares							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	85	2	9	3	35	134
	%	25.60%	1.03%	13.63%	5.66%	29.66%	17.58%
- Yes	N	3	--	1	--	1	5
	%	0.90%	--	1.51%	--	0.84%	0.65%
Reason for stopping Aricept - Nausea							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	82	2	9	3	32	128
	%	24.69%	1.03%	13.63%	5.66%	27.11%	16.79%
- Yes	N	6	--	1	--	4	11
	%	1.80%	--	1.51%	--	3.38%	1.44%
Reason for stopping Aricept - Weight Loss							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	88	2	9	3	36	138
	%	26.50%	1.03%	13.63%	5.66%	30.50%	18.11%
- Yes	N	--	--	1	--	--	1
	%	--	--	1.51%	--	--	0.13%
Reason for stopping Aricept - Vomiting							
	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	86	2	10	3	35	136
	%	25.90%	1.03%	15.15%	5.66%	29.66%	17.84%
- Yes	N	2	--	--	--	1	3
	%	0.60%	--	--	--	0.84%	0.39%
Reason for stopping Aricept - Other							

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Aricept - Other							
-	N	244	191	56	50	82	623
	%	73.49%	98.96%	84.84%	94.33%	69.49%	81.75%
- No	N	57	1	6	2	23	89
	%	17.16%	0.51%	9.09%	3.77%	19.49%	11.67%
- Yes	N	31	1	4	1	13	50
	%	9.33%	0.51%	6.06%	1.88%	11.01%	6.56%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Amount of time Aricept was used							
	N	273	191	56	50	82	652
	%	82.22%	98.96%	84.84%	94.33%	69.49%	85.56%
- < 1 Month	N	12	--	2	--	2	16
	%	3.61%	--	3.03%	--	1.69%	2.09%
- 1-2 Months	N	5	--	3	1	4	13
	%	1.50%	--	4.54%	1.88%	3.38%	1.70%
- 3-6 Months	N	12	1	1	--	7	21
	%	3.61%	0.51%	1.51%	--	5.93%	2.75%
- 7-12 Months	N	6	--	--	1	2	9
	%	1.80%	--	--	1.88%	1.69%	1.18%
- > 12 Months	N	24	1	4	1	21	51
	%	7.22%	0.51%	6.06%	1.88%	17.79%	6.69%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject has ever taken Exelon tablets (MTH Q2)							
	N	3	2	--	--	8	13
	%	0.90%	1.03%	--	--	6.77%	1.70%
- No	N	287	190	65	53	97	692
	%	86.44%	98.44%	98.48%	100.00%	82.20%	90.81%
- Yes	N	42	1	1	--	13	57
	%	12.65%	0.51%	1.51%	--	11.01%	7.48%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject still taking Exelon tablets (MTH Q2A)							
	N	290	192	65	53	105	705
	%	87.34%	99.48%	98.48%	100.00%	88.98%	92.51%

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Subject still taking Exelon tablets (MTH Q2A) - No	N	29	1	--	--	9	39
	%	8.73%	0.51%	--	--	7.62%	5.11%
- Yes	N	13	--	1	--	4	18
	%	3.91%	--	1.51%	--	3.38%	2.36%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
		332	193	66	53	118
Reason for stopping Exelon tablets - Did not help						
	N	303	192	66	53	109
	%	91.26%	99.48%	100.00%	100.00%	92.37%
- No	N	22	1	--	--	8
	%	6.62%	0.51%	--	--	6.77%
- Yes	N	7	--	--	--	1
	%	2.10%	--	--	--	0.84%
Reason for stopping Exelon tablets - Abdominal Pain/Cramps						
	N	303	192	66	53	109
	%	91.26%	99.48%	100.00%	100.00%	92.37%
- No	N	27	1	--	--	7
	%	8.13%	0.51%	--	--	5.93%
- Yes	N	2	--	--	--	2
	%	0.60%	--	--	--	1.69%
Reason for stopping Exelon tablets - Agitation/Restlessness						
	N	303	192	66	53	109
	%	91.26%	99.48%	100.00%	100.00%	92.37%
- No	N	26	1	--	--	8
	%	7.83%	0.51%	--	--	6.77%
- Yes	N	3	--	--	--	1
	%	0.90%	--	--	--	0.84%
Reason for stopping Exelon tablets - Anorexia						
	N	303	192	66	53	109
	%	91.26%	99.48%	100.00%	100.00%	92.37%
- No	N	29	1	--	--	9
	%	8.73%	0.51%	--	--	7.62%
Reason for stopping Exelon tablets - Confusion						
	N	303	192	66	53	109
	%	91.26%	99.48%	100.00%	100.00%	92.37%

(Continued)

Medication Treatment History		All
		762
Reason for stopping Exelon tablets - Did not help		
- .	N	723
	%	94.88%
- No	N	31
	%	4.06%
- Yes	N	8
	%	1.04%
Reason for stopping Exelon tablets - Abdominal Pain/Cramps		
- .	N	723
	%	94.88%
- No	N	35
	%	4.59%
- Yes	N	4
	%	0.52%
Reason for stopping Exelon tablets - Agitation/Restlessness		
- .	N	723
	%	94.88%
- No	N	35
	%	4.59%
- Yes	N	4
	%	0.52%
Reason for stopping Exelon tablets - Anorexia		
- .	N	723
	%	94.88%
- No	N	39
	%	5.11%
Reason for stopping Exelon tablets - Confusion		
- .	N	723
	%	94.88%

(Continued)

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
Reason for stopping Exelon tablets - Confusion						
- No	N	29	1	--	--	9
	%	8.73%	0.51%	--	--	7.62%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		All
Reason for stopping Exelon tablets - Confusion		762
- No	N	39
	%	5.11%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Exelon tablets - Diarrhea							
	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	26	--	--	--	8	34
	%	7.83%	--	--	--	6.77%	4.46%
- Yes	N	3	1	--	--	1	5
	%	0.90%	0.51%	--	--	0.84%	0.65%
Reason for stopping Exelon tablets - Dizziness							
	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	28	1	--	--	9	38
	%	8.43%	0.51%	--	--	7.62%	4.98%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Exelon tablets - Fatigue							
	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	29	1	--	--	8	38
	%	8.73%	0.51%	--	--	6.77%	4.98%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Exelon tablets - Headache							
	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	29	1	--	--	8	38
	%	8.73%	0.51%	--	--	6.77%	4.98%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Exelon tablets - Insomnia							

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Exelon tablets - Insomnia							
-	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	29	1	--	--	9	39
	%	8.73%	0.51%	--	--	7.62%	5.11%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Exelon tablets - Nightmares							
-	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	29	1	--	--	9	39
	%	8.73%	0.51%	--	--	7.62%	5.11%
Reason for stopping Exelon tablets - Nausea							
-	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	24	1	--	--	7	32
	%	7.22%	0.51%	--	--	5.93%	4.19%
- Yes	N	5	--	--	--	2	7
	%	1.50%	--	--	--	1.69%	0.91%
Reason for stopping Exelon tablets - Weight Loss							
-	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	28	1	--	--	9	38
	%	8.43%	0.51%	--	--	7.62%	4.98%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Exelon tablets - Vomiting							
-	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%
- No	N	27	1	--	--	8	36
	%	8.13%	0.51%	--	--	6.77%	4.72%
- Yes	N	2	--	--	--	1	3
	%	0.60%	--	--	--	0.84%	0.39%
Reason for stopping Exelon tablets - Other							
-	N	303	192	66	53	109	723
	%	91.26%	99.48%	100.00%	100.00%	92.37%	94.88%

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Exelon tablets - Other							
- No	N	18	1	--	--	6	25
	%	5.42%	0.51%	--	--	5.08%	3.28%
- Yes	N	11	--	--	--	3	14
	%	3.31%	--	--	--	2.54%	1.83%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Amount of time Exelon tablets was used							
	N	314	192	66	53	109	734
	%	94.57%	99.48%	100.00%	100.00%	92.37%	96.32%
- < 1 Month	N	5	--	--	--	1	6
	%	1.50%	--	--	--	0.84%	0.78%
- 1-2 Months	N	3	--	--	--	--	3
	%	0.90%	--	--	--	--	0.39%
- 3-6 Months	N	5	1	--	--	2	8
	%	1.50%	0.51%	--	--	1.69%	1.04%
- 7-12 Months	N	1	--	--	--	1	2
	%	0.30%	--	--	--	0.84%	0.26%
- > 12 Months	N	4	--	--	--	5	9
	%	1.20%	--	--	--	4.23%	1.18%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject has ever used Exelon Patches (MTH Q3)							
	N	138	2	--	--	8	148
	%	41.56%	1.03%	--	--	6.77%	19.42%
- No	N	164	190	60	52	80	546
	%	49.39%	98.44%	90.90%	98.11%	67.79%	71.65%
- Yes	N	30	1	6	1	30	68
	%	9.03%	0.51%	9.09%	1.88%	25.42%	8.92%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject still using Exelon Patches (MTH Q3A)							
	N	302	192	60	52	88	694
	%	90.96%	99.48%	90.90%	98.11%	74.57%	91.07%

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Subject still using Exelon Patches (MTH Q3A) - No	N	17	1	2	1	18	39
	%	5.12%	0.51%	3.03%	1.88%	15.25%	5.11%
- Yes	N	13	--	4	--	12	29
	%	3.91%	--	6.06%	--	10.16%	3.80%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Exelon Patches - Did not help							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	12	1	1	1	13	28
	%	3.61%	0.51%	1.51%	1.88%	11.01%	3.67%
- Yes	N	5	--	1	--	5	11
	%	1.50%	--	1.51%	--	4.23%	1.44%
Reason for stopping Exelon Patches - Abdominal Pain/Cramps							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	17	1	2	1	17	38
	%	5.12%	0.51%	3.03%	1.88%	14.40%	4.98%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Exelon Patches - Agitation/Restlessness							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	16	1	2	1	18	38
	%	4.81%	0.51%	3.03%	1.88%	15.25%	4.98%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Exelon Patches - Anorexia							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	17	1	2	1	18	39
	%	5.12%	0.51%	3.03%	1.88%	15.25%	5.11%
Reason for stopping Exelon Patches - Confusion							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	15	1	2	1	18	37

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Exelon Patches - Confusion							
- No	%	4.51%	0.51%	3.03%	1.88%	15.25%	4.85%
- Yes	N	2	--	--	--	--	2
	%	0.60%	--	--	--	--	0.26%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Exelon Patches - Diarrhea							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	16	--	2	1	18	37
	%	4.81%	--	3.03%	1.88%	15.25%	4.85%
- Yes	N	1	1	--	--	--	2
	%	0.30%	0.51%	--	--	--	0.26%
Reason for stopping Exelon Patches - Dizziness							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	16	1	2	1	18	38
	%	4.81%	0.51%	3.03%	1.88%	15.25%	4.98%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Exelon Patches - Fatigue							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	17	1	2	1	17	38
	%	5.12%	0.51%	3.03%	1.88%	14.40%	4.98%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Exelon Patches - Headache							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	17	1	2	1	18	39
	%	5.12%	0.51%	3.03%	1.88%	15.25%	5.11%
Reason for stopping Exelon Patches - Insomnia							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	17	1	2	1	18	39
	%	5.12%	0.51%	3.03%	1.88%	15.25%	5.11%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Exelon Patches - Nightmares							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	17	1	2	1	18	39
	%	5.12%	0.51%	3.03%	1.88%	15.25%	5.11%
Reason for stopping Exelon Patches - Nausea							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	16	1	2	1	17	37
	%	4.81%	0.51%	3.03%	1.88%	14.40%	4.85%
- Yes	N	1	--	--	--	1	2
	%	0.30%	--	--	--	0.84%	0.26%
Reason for stopping Exelon Patches - Weight Loss							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	16	1	2	1	18	38
	%	4.81%	0.51%	3.03%	1.88%	15.25%	4.98%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Exelon Patches - Vomiting							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- No	N	16	1	2	1	17	37
	%	4.81%	0.51%	3.03%	1.88%	14.40%	4.85%
- Yes	N	1	--	--	--	1	2
	%	0.30%	--	--	--	0.84%	0.26%
Reason for stopping Exelon Patches - Other							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Exelon Patches - Other							
- No	N	7	1	1	--	6	15
	%	2.10%	0.51%	1.51%	--	5.08%	1.96%
- Yes	N	10	--	1	1	12	24
	%	3.01%	--	1.51%	1.88%	10.16%	3.14%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Amount of time Exelon Patches was used							
	N	315	192	64	52	100	723
	%	94.87%	99.48%	96.96%	98.11%	84.74%	94.88%
- < 1 Month	N	4	--	--	--	2	6
	%	1.20%	--	--	--	1.69%	0.78%
- 1-2 Months	N	2	1	--	--	2	5
	%	0.60%	0.51%	--	--	1.69%	0.65%
- 3-6 Months	N	6	--	1	--	2	9
	%	1.80%	--	1.51%	--	1.69%	1.18%
- 7-12 Months	N	--	--	--	--	4	4
	%	--	--	--	--	3.38%	0.52%
- > 12 Months	N	5	--	1	1	8	15
	%	1.50%	--	1.51%	1.88%	6.77%	1.96%

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
		332	193	66	53	118
Subject has ever taken Razadyne/Reminyl (MTH Q4)						
-	N	3	2	--	--	8
	%	0.90%	1.03%	--	--	6.77%
- No	N	294	191	65	53	96
	%	88.55%	98.96%	98.48%	100.00%	81.35%
- Yes	N	35	--	1	--	14
	%	10.54%	--	1.51%	--	11.86%

Medication Treatment History		All
		762
Subject has ever taken Razadyne/Reminyl (MTH Q4)		
-	N	13
	%	1.70%
- No	N	699
	%	91.73%
- Yes	N	50
	%	6.56%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject still taking Razadyne/Reminyl (MTH Q4A)							
-	N	297	193	65	53	104	712
	%	89.45%	100.00%	98.48%	100.00%	88.13%	93.43%
- No	N	18	--	1	--	9	28
	%	5.42%	--	1.51%	--	7.62%	3.67%
- Yes	N	17	--	--	--	5	22
	%	5.12%	--	--	--	4.23%	2.88%

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
		332	193	66	53	118
Reason for stopping Exelon Patches - Did not help						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	9	--	1	--	8
	%	2.71%	--	1.51%	--	6.77%
- Yes	N	9	--	--	--	1
	%	2.71%	--	--	--	0.84%
Reason for stopping Exelon Patches - Abdominal Pain/Cramps						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	18	--	1	--	8
	%	5.42%	--	1.51%	--	6.77%
- Yes	N	--	--	--	--	1
	%	--	--	--	--	0.84%
Reason for stopping Exelon Patches - Agitation/Restlessness						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	14	--	1	--	7
	%	4.21%	--	1.51%	--	5.93%
- Yes	N	4	--	--	--	2
	%	1.20%	--	--	--	1.69%
Reason for stopping Exelon Patches - Anorexia						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	18	--	1	--	9
	%	5.42%	--	1.51%	--	7.62%
Reason for stopping Exelon Patches - Confusion						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	17	--	1	--	9

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		All
		762
Reason for stopping Exelon Patches - Did not help		
- .	N	734
	%	96.32%
- No	N	18
	%	2.36%
- Yes	N	10
	%	1.31%
Reason for stopping Exelon Patches - Abdominal Pain/Cramps		
- .	N	734
	%	96.32%
- No	N	27
	%	3.54%
- Yes	N	1
	%	0.13%
Reason for stopping Exelon Patches - Agitation/Restlessness		
- .	N	734
	%	96.32%
- No	N	22
	%	2.88%
- Yes	N	6
	%	0.78%
Reason for stopping Exelon Patches - Anorexia		
- .	N	734
	%	96.32%
- No	N	28
	%	3.67%
Reason for stopping Exelon Patches - Confusion		
- .	N	734
	%	96.32%
- No	N	27

(Continued)

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
Reason for stopping Exelon Patches - Confusion						
- No	%	5.12%	--	1.51%	--	7.62%
- Yes	N	1	--	--	--	--
	%	0.30%	--	--	--	--

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		All
Reason for stopping Exelon Patches - Confusion		762
- No	%	3.54%
- Yes	N	1
	%	0.13%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Exelon Patches - Diarrhea							
	N	314	193	65	53	109	734
	%	94.57%	100.00%	98.48%	100.00%	92.37%	96.32%
- No	N	18	--	1	--	9	28
	%	5.42%	--	1.51%	--	7.62%	3.67%
Reason for stopping Exelon Patches - Dizziness							
	N	314	193	65	53	109	734
	%	94.57%	100.00%	98.48%	100.00%	92.37%	96.32%
- No	N	18	--	1	--	9	28
	%	5.42%	--	1.51%	--	7.62%	3.67%
Reason for stopping Exelon Patches - Fatigue							
	N	314	193	65	53	109	734
	%	94.57%	100.00%	98.48%	100.00%	92.37%	96.32%
- No	N	18	--	1	--	8	27
	%	5.42%	--	1.51%	--	6.77%	3.54%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Exelon Patches - Headache							
	N	314	193	65	53	109	734
	%	94.57%	100.00%	98.48%	100.00%	92.37%	96.32%
- No	N	18	--	1	--	8	27
	%	5.42%	--	1.51%	--	6.77%	3.54%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Exelon Patches - Insomnia							
	N	314	193	65	53	109	734
	%	94.57%	100.00%	98.48%	100.00%	92.37%	96.32%
- No	N	18	--	1	--	9	28
	%	5.42%	--	1.51%	--	7.62%	3.67%

Medication Treatment History		Status				
		.	Normal	MCI	QCI	Dementia
		332	193	66	53	118
Reason for stopping Exelon Patches - Nightmares						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	17	--	1	--	9
	%	5.12%	--	1.51%	--	7.62%
- Yes	N	1	--	--	--	--
	%	0.30%	--	--	--	--
Reason for stopping Exelon Patches - Nausea						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	18	--	1	--	9
	%	5.42%	--	1.51%	--	7.62%
Reason for stopping Exelon Patches - Weight Loss						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	18	--	1	--	9
	%	5.42%	--	1.51%	--	7.62%
Reason for stopping Exelon Patches - Vomiting						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	18	--	1	--	9
	%	5.42%	--	1.51%	--	7.62%
Reason for stopping Exelon Patches - Other						
	N	314	193	65	53	109
	%	94.57%	100.00%	98.48%	100.00%	92.37%
- No	N	13	--	--	--	4
	%	3.91%	--	--	--	3.38%
- Yes	N	5	--	1	--	5
	%	1.50%	--	1.51%	--	4.23%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		All
		762
Reason for stopping Exelon Patches - Nightmares		
-	N	734
	%	96.32%
- No	N	27
	%	3.54%
- Yes	N	1
	%	0.13%
Reason for stopping Exelon Patches - Nausea		
-	N	734
	%	96.32%
- No	N	28
	%	3.67%
Reason for stopping Exelon Patches - Weight Loss		
-	N	734
	%	96.32%
- No	N	28
	%	3.67%
Reason for stopping Exelon Patches - Vomiting		
-	N	734
	%	96.32%
- No	N	28
	%	3.67%
Reason for stopping Exelon Patches - Other		
-	N	734
	%	96.32%
- No	N	17
	%	2.23%
- Yes	N	11
	%	1.44%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Amount of time Exelon Patches was used							
	N	321	193	65	53	109	741
	%	96.68%	100.00%	98.48%	100.00%	92.37%	97.24%
- < 1 Month	N	1	--	--	--	3	4
	%	0.30%	--	--	--	2.54%	0.52%
- 1-2 Months	N	--	--	1	--	1	2
	%	--	--	1.51%	--	0.84%	0.26%
- 3-6 Months	N	4	--	--	--	1	5
	%	1.20%	--	--	--	0.84%	0.65%
- 7-12 Months	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
- > 12 Months	N	5	--	--	--	4	9
	%	1.50%	--	--	--	3.38%	1.18%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject has ever taken Namenda (MTH Q5)							
	N	3	2	--	--	8	13
	%	0.90%	1.03%	--	--	6.77%	1.70%
- No	N	200	191	54	49	34	528
	%	60.24%	98.96%	81.81%	92.45%	28.81%	69.29%
- Yes	N	129	--	12	4	76	221
	%	38.85%	--	18.18%	7.54%	64.40%	29.00%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Subject still taking Namenda (MTH Q5A)							
	N	203	193	54	49	42	541
	%	61.14%	100.00%	81.81%	92.45%	35.59%	70.99%

(Continued)

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Subject still taking Namenda (MTH Q5A) - No	N	47	--	1	4	30	82
	%	14.15%	--	1.51%	7.54%	25.42%	10.76%
- Yes	N	82	--	11	--	46	139
	%	24.69%	--	16.66%	--	38.98%	18.24%

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Namenda - Did not help							
	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	21	--	--	3	21	45
	%	6.32%	--	--	5.66%	17.79%	5.90%
- Yes	N	26	--	1	1	9	37
	%	7.83%	--	1.51%	1.88%	7.62%	4.85%
Reason for stopping Namenda - Abdominal Pain/Cramps							
	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	46	--	1	4	30	81
	%	13.85%	--	1.51%	7.54%	25.42%	10.62%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Namenda - Agitation/Restlessness							
	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	45	--	1	4	29	79
	%	13.55%	--	1.51%	7.54%	24.57%	10.36%
- Yes	N	2	--	--	--	1	3
	%	0.60%	--	--	--	0.84%	0.39%
Reason for stopping Namenda - Anorexia							
	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	47	--	1	4	30	82
	%	14.15%	--	1.51%	7.54%	25.42%	10.76%
Reason for stopping Namenda - Confusion							
	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	46	--	1	4	28	79

(Continued)

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
Reason for stopping Namenda - Confusion							
- No	%	13.85%	--	1.51%	7.54%	23.72%	10.36%
- Yes	N	1	--	--	--	2	3
	%	0.30%	--	--	--	1.69%	0.39%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Namenda - Diarrhea							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	44	--	1	4	29	78
	%	13.25%	--	1.51%	7.54%	24.57%	10.23%
- Yes	N	3	--	--	--	1	4
	%	0.90%	--	--	--	0.84%	0.52%
Reason for stopping Namenda - Dizziness							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	45	--	1	4	29	79
	%	13.55%	--	1.51%	7.54%	24.57%	10.36%
- Yes	N	2	--	--	--	1	3
	%	0.60%	--	--	--	0.84%	0.39%
Reason for stopping Namenda - Fatigue							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	46	--	1	4	30	81
	%	13.85%	--	1.51%	7.54%	25.42%	10.62%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Namenda - Headache							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	47	--	1	4	30	82
	%	14.15%	--	1.51%	7.54%	25.42%	10.76%
Reason for stopping Namenda - Insomnia							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	47	--	1	4	30	82
	%	14.15%	--	1.51%	7.54%	25.42%	10.76%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Reason for stopping Namenda - Nightmares							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	47	--	1	4	30	82
	%	14.15%	--	1.51%	7.54%	25.42%	10.76%
Reason for stopping Namenda - Nausea							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	46	--	1	4	30	81
	%	13.85%	--	1.51%	7.54%	25.42%	10.62%
- Yes	N	1	--	--	--	--	1
	%	0.30%	--	--	--	--	0.13%
Reason for stopping Namenda - Weight Loss							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	47	--	1	4	30	82
	%	14.15%	--	1.51%	7.54%	25.42%	10.76%
Reason for stopping Namenda - Vomiting							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	47	--	1	4	29	81
	%	14.15%	--	1.51%	7.54%	24.57%	10.62%
- Yes	N	--	--	--	--	1	1
	%	--	--	--	--	0.84%	0.13%
Reason for stopping Namenda - Other							
- .	N	285	193	65	49	88	680
	%	85.84%	100.00%	98.48%	92.45%	74.57%	89.23%
- No	N	33	--	1	2	12	48
	%	9.93%	--	1.51%	3.77%	10.16%	6.29%
- Yes	N	14	--	--	2	18	34
	%	4.21%	--	--	3.77%	15.25%	4.46%

Note: When multiple observations are taken for a patient, only latest visit is accounted for in summary.

Medication Treatment History		Status					All
		.	Normal	MCI	QCI	Dementia	
		332	193	66	53	118	762
Amount of time Namenda was used							
	N	302	193	65	49	88	697
	%	90.96%	100.00%	98.48%	92.45%	74.57%	91.46%
- < 1 Month	N	3	--	1	--	1	5
	%	0.90%	--	1.51%	--	0.84%	0.65%
- 1-2 Months	N	2	--	--	1	5	8
	%	0.60%	--	--	1.88%	4.23%	1.04%
- 3-6 Months	N	5	--	--	1	4	10
	%	1.50%	--	--	1.88%	3.38%	1.31%
- 7-12 Months	N	2	--	--	--	5	7
	%	0.60%	--	--	--	4.23%	0.91%
- > 12 Months	N	18	--	--	2	15	35
	%	5.42%	--	--	3.77%	12.71%	4.59%